



碧云天生物技术/Beyotime Biotechnology
 订货热线: 400-1683301或800-8283301
 订货e-mail: order@beyotime.com
 技术咨询: info@beyotime.com
 网址: http://www.beyotime.com

Bcl-2 Rabbit Polyclonal Antibody

产品编号	产品名称	包装
AF0060	Bcl-2 Rabbit Polyclonal Antibody	50μl

产品简介:

来源	用途	交叉反应性	分子量
Rabbit	WB, FC, IHC, ICC, IF	H	26KDa

WB, Western blot; IP, Immunoprecipitation; IF, Immunofluorescence; IHC, Immunohistochemistry; ICC, Immunocytochemistry; FC, Flow Cytometry; ELISA, Enzyme-linked Immunosorbent Assay; ChIP, Chromatin Immunoprecipitation Assay.

H, Human; M, Mouse; R, Rat; C, Chicken; Cw, Cow; Dg, Dog; Gp, Guinea pig; Hm, Hamster; Hr, Horse; Mk, Monkey; Pg, Pig; Rb, Rabbit; S, Sheep; Z, Zebrafish; All, all species expected.

- 配套提供了Western一抗稀释液, 可以用于Western检测或其它适当用途时的一抗稀释。
- 建议抗体使用时的稀释比例如下(实际使用时需根据抗原水平的高低作适当调整):

WB	IP	IF	IHC	ICC	FC	ELISA	ChIP
1:500-1:1000	-	1:200	1:200	1:200	1:50-1:100	-	-

- 抗体详细信息如下:

About this Antibody	
Name	Bcl-2 Rabbit Polyclonal Antibody
Category	Polyclonal antibody(pAb); Primary antibody
Isotype	IgG
Purification	Peptide affinity purified
About the Immunogen	
Immunogen	This antibody is produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to human Bcl-2.
Gene ID	596
SwissProt	P10415
Synonyms	Bcl-2; PPP1R50
Category	Apoptosis
Background	Damage to the Bcl-2 gene has been identified as a cause of a number of cancers, including melanoma, breast, prostate, chronic lymphocytic leukemia, and lung cancer, and a possible cause of schizophrenia and autoimmunity. It is also a cause of resistance to cancer treatments. Antibodies to Bcl-2 can be used with immunohistochemistry to identify cells containing the antigen. In healthy tissue, these antibodies will react with B-cells in the mantle zone, as well as some T-cells. However, there is a considerable increase in positive cells in follicular lymphoma, as well as many other forms of cancer. In some cases, the presence or absence of Bcl-2 staining in biopsies may be significant for the patient's prognosis or likelihood of relapse.

包装清单:

产品编号	产品名称	包装
AF0060	Bcl-2 Rabbit Polyclonal Antibody	50μl
AZ050	Western一抗稀释液	50ml
-	说明书	1份

保存条件:

Bcl-2 Rabbit Polyclonal Antibody -20°C保存, Western一抗稀释液-20°C或4°C保存, 一年有效。Western一抗稀释液优先推荐4°C保存, 长期不使用可以考虑-20°C保存, 但冻融可能会导致出现轻微的浑浊和少量不溶物。

注意事项:

- 如果本抗体用于Western blot (WB)、免疫荧光(IF)、免疫细胞化学(ICC)等实验, 请注意回收使用过的稀释抗体。回收的抗体通常至少可以重复使用5-10次。稀释后的抗体, 包括已经使用过的稀释抗体, 请4°C保存。
- 回收后重复使用的抗体, 使用方法同新鲜稀释的抗体。如果在重复使用过程中发现抗体出现轻微混浊现象, 可以10,000g离心1-3分钟, 取上清用于后续检测。如果回收的抗体出现明显的絮状物或长霉长菌等情况, 则可以考虑废弃该抗体。
- 提供的Western一抗稀释液也可以用于免疫荧光(IF)、免疫组化(IHC)、免疫细胞化学(ICC)等适当用途。如果希望获得最佳的检测效果, 请考虑使用上述检测专用的一抗稀释液。
- 本产品仅限于专业人员的科学研究用, 不得用于临床诊断或治疗, 不得用于食品或药品, 不得存放于普通住宅内。
- 为了您的安全和健康, 请穿实验服并戴一次性手套操作。

使用说明:

请根据抗体的实际用途选择相应的使用方法。

1. Western检测:

- 按照推荐的稀释比例用碧云天提供的Western一抗稀释液稀释抗体。
- 把经过封闭的蛋白膜与稀释好的一抗4°C缓慢摇动过夜或室温缓慢摇动2小时, 确保稀释的抗体至少能在摇动的瞬间覆盖蛋白膜。
- 回收稀释的一抗, 4°C保存以备下次继续使用。
- 按照Western的实验步骤进行后续的洗涤、二抗孵育、洗涤和检测等操作。具体操作可以参考如下网页:
<http://www.beyotime.com/support/western.htm>

2. 免疫染色:

可以使用碧云天生产的免疫染色一抗稀释液(P0103)稀释抗体, 使用后注意回收稀释好的一抗, 具体操作可以参考如下网页: <http://www.beyotime.com/support/immunol-staining.htm>

3. 其它实验操作请自行参考适当的protocol进行。

4. 代表性图片:




Fig. 1. Western blot analysis of Bcl-2 on different cell lysates using anti-Bcl-2 antibody at 1/500 dilution. Positive control: Lane1: HL-60 Lane2: Hela Lane3: HepG2 Lane 4:A431




Fig. 2. Immunocytochemistry staining Bcl-2 in HeLa cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig. 3. Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue using anti-Bcl-2 rabbit polyclonal antibody. Counter stained with hematoxylin.




Fig. 4. Flow cytometric analysis of HeLa cells with Bcl-2 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Goat anti rabbit IgG (FITC) was

used as the secondary antibody.

相关产品:

产品编号	产品名称	包装
P0006	Bradford蛋白浓度测定试剂盒	1000次
P0010	BCA蛋白浓度测定试剂盒(增强型)	500次
P0012	BCA蛋白浓度测定试剂盒	500次
P0012A	SDS-PAGE凝胶配制试剂盒	1盒
P0012AC	SDS-PAGE凝胶快速配制试剂盒	1盒
P0013	Western及IP细胞裂解液	100ml
P0013B	RIPA裂解液(强)	100ml
P0014B	SDS-PAGE电泳液	10×1L
P0015	SDS-PAGE蛋白上样缓冲液(5X)	2ml
P0018	BeyoECL Plus(超敏ECL化学发光试剂盒)	共100ml
P0018A	BeyoECL Star(特超敏ECL化学发光试剂盒)	共100ml
P0020	显影定影试剂盒	各1升
P0021B	Western转膜液	10×1L
P0023A	Western一抗稀释液	100ml
P0023B	Western封闭液	100ml
P0023C6	Western洗涤液(10X)	10×100ml
P0023D	Western二抗稀释液	100ml
P0025	Western一抗二抗去除液	250ml
P0252	QuickBlock™ Western封闭液	100ml
P0256	QuickBlock™ Western一抗稀释液	100ml
P0258	QuickBlock™ Western二抗稀释液	100ml
P0066	预染蛋白质分子量标准	200μl
P0068	彩色预染蛋白质分子量标准	200μl
P0071	BeyoColor™彩色预染蛋白质分子量标准	200μl
P0098	免疫染色固定液	100ml
P0102	免疫染色封闭液	100ml
P0103	免疫染色一抗稀释液	100ml
P0106L	免疫染色洗涤液(10X)	10×100ml
P0108	免疫荧光染色二抗稀释液	100ml
P0110	免疫染色(非荧光)二抗稀释液	100ml
P0260	QuickBlock™免疫染色封闭液	100ml
P0262	QuickBlock™免疫染色一抗稀释液	100ml
P0265	QuickBlock™免疫荧光染色二抗稀释液	100ml
P0267	QuickBlock™免疫组化染色二抗稀释液	100ml
P0126	抗荧光淬灭封片液	5ml
FFP24	PVDF膜(进口分装, 6.6×8.5cm, 0.2μm)	20张/包装
FFP32	PVDF膜(进口原装, 6.6×8.5cm, 0.45μm)	20张/包装
FFP51	转印滤纸(7.5×10cm)	100片/包装
FF057	X-OMAT BT胶片(原柯达, 5×7英寸)	100张/盒
FFC58	压片暗盒(5×7英寸)	1个

使用本产品的文献:

1. Meng Z, Liu Y, Wang J, Fan H, Fang H, Li S, Yuan L, Liu C, Peng Y, Zhao W, Wang L, Li J, Feng J . Histone demethylase KDM7A is required for stem cell maintenance and apoptosis inhibition in breast cancer. J Cell Physiol. 2020 Feb 235(2):932-943.
2. Liu P, Dong J. . Protective effects of carnolic acid against mitochondria-mediated injury in H9c2 cardiomyocytes induced by hypoxia/reoxygenation. Exp Ther Med. 2017 Dec;14(6):5629-5634.
3. Gao H, Zhang Y, Dong L, Qu XY, Tao LN, Zhang YM, Zhai JH, Song YQ . Triptolide induces autophagy and apoptosis through ERK activation in human breast cancer MCF-7 cells. Exp Ther Med. 2018 Apr;15(4):3413-3419.
4. Wang C, Ke H, Xu X, Chen J, Sun D, Ji F . Protective effect of nicorandil on collapse-induced lung injury in rabbits by inhibiting apoptosis. Int J Mol Med. 2019 Aug 44(2):725-736.

5. Li SS, Jiang WL, Xiao WQ, Li K, Zhang YF, Guo XY, Dai YQ, Zhao QY, Jiang MJ, Lu ZJ, Wan R . KMT2D deficiency enhances the anti-cancer activity of L48H37 in pancreatic ductal adenocarcinoma. *World J Gastrointest Oncol*. 2019 Aug 15 11(8):599-621.
6. Yu Y, Lang XY, Li XX, Gu RZ, Liu QS, Lan R, Qin XY . 2,3,5,4'-Tetrahydroxystilbene-2-O- β -d-glucoside attenuates MPP+/MPTP-induced neurotoxicity in vitro and in vivo by restoring the BDNF-TrkB and FGF2-Akt signaling axis and inhibition of apoptosis. *Food Funct*. 2019 Sep 1 10(9):6009-6019.
7. Jin L, Wang Q, Chen J, Wang Z, Xin H, Zhang D . Efficient Delivery of Therapeutic siRNA by Fe₃O₄ Magnetic Nanoparticles into Oral Cancer Cells. *Pharmaceutics*. 2019 Nov 17 11(11). pii: E615.
8. Yi Yang, Xiaoyu Luo, Mayila Yasheng, Jun Zhao, Jinyu Li, Jinyao Li . Ergosterol peroxide from *Pleurotus ferulae* inhibits gastrointestinal tumor cell growth through induction of apoptosis via reactive oxygen species and endoplasmic reticulum stress *Food Funct*. 2020 May 1;11(5):4171-4184.
9. Junyang Zhou, Zhixiao Wang, Yun He, Xinxia Luo, Wenjun Zhang, Li Yu, Xiuying Chen, Xiju He, Yahong Yuan, Xiaoli Wang, Xinrong Guo, Junming Tang, Mangan Zhu, Dongsheng Li, Yan Ding . Qiliqiangxin reduced cardiomyocytes apoptosis and improved heart function in infarcted heart through Pink1/Parkin -mediated mitochondrial autophagy *BMC Complement Med Ther*. 2020 Jul 2;20(1):203.
10. Teng-Teng Ren, Sheng-Rui Fan, Xiu-Yuan Lang, Yun Yu, Rongfeng Lan, Xiao-Yan Qin . 2,3,5,4'-tetrahydroxystilbene-2-O- β -D-glucoside eliminates staurosporine-induced cytotoxicity by restoring BDNF-TrkB/Akt signaling axis *Int J Med Sci*. 2020 Aug 19;17(14):2207-2213.
11. Jun Hui Xu, Yong Jun Guan, Yi Chao Zhang, Zhen Dong Qiu, Yu Zhou, Chen Chen, Jia Yu, Wei Xing Wang . ADAM15 correlates with prognosis, immune infiltration and apoptosis in hepatocellular carcinoma *Aging (Albany NY)*. 2021 Aug 23;13(16):20395-20417.
12. Jian Zhao, Hwei Li, Wanying Li, Zixian Wang, Zhewen Dong, Huan Lan, Chengqiang Wang, Jia-Le Song . Effects of Sinapic Acid Combined with Cisplatin on the Apoptosis and Autophagy of the Hepatoma Cells HepG2 and SMMC-7721 *Evid Based Complement Alternat Med*. 2021 Oct 12;2021:6095963.
13. Xierenguli Halike, Jinyu Li, Pengfei Yuan, Kaimeiliya Yasheng, Min Chen, Lijie Xia, Jinyao Li . The petroleum ether extract of *Brassica rapa* L. induces apoptosis of lung adenocarcinoma cells via the mitochondria-dependent pathway *Food Funct*. 2021 Oct 19;12(20):10023-10039.

Version 2024.03.12