



碧云天生物技术/Beyotime Biotechnology
订货热线: 400-1683301或800-8283301
订货e-mail: order@beyotime.com
技术咨询: info@beyotime.com
网址: http://www.beyotime.com

NF-κB p65 Rabbit Polyclonal Antibody

产品编号	产品名称	包装
AF0246	NF-κB p65 Rabbit Polyclonal Antibody	50μl

产品简介:

来源	用途	交叉反应性	分子量
Rabbit	WB, FC, IHC	H, M, R	65KDa

WB, Western blot; IP, Immunoprecipitation; IF, Immunofluorescence; IHC, Immunohistochemistry; ICC, Immunocytochemistry; FC, Flow Cytometry; ELISA, Enzyme-linked Immunosorbent Assay; ChIP, Chromatin Immunoprecipitation Assay.

H, Human; M, Mouse; R, Rat; C, Chicken; Cw, Cow; Dg, Dog; Gp, Guinea pig; Hm, Hamster; Hr, Horse; Mk, Monkey; Pg, Pig; Rb, Rabbit; S, Sheep; Z, Zebrafish; All, all species expected.

➤ 配套提供了Western一抗稀释液，可以用于Western检测或其它适当用途时的一抗稀释。

➤ 建议抗体使用时的稀释比如下(实际使用时需根据抗原水平的高低作适当调整):

WB	IP	IF	IHC	ICC	FC	ELISA	ChIP
1:1000-1:2000	-	-	1:100-1:200	-	1:50-1:100	-	-

抗体详细信息如下:

About this Antibody	
Name	NF-κB p65 Rabbit Polyclonal Antibody
Category	Polyclonal antibody(pAb); Primary antibody
Isotype	IgG
Purification	Peptide affinity purified
About the Immunogen	
Immunogen	This antibody is produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to N-terminal NF-κB p65.
Gene ID	5970
SwissProt	Q04206
Synonyms	NFKB3; Nuclear factor NF-kappa-B p65 subunit; RELA; TF65; Transcription factor p65; p65
Category	Immunology and Inflammation
Background	NF-κB is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-κB is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. In unstimulated cells, NF-κB is sequestered in the cytoplasm by IκB inhibitory proteins. NF-κB-activating agents can induce the phosphorylation of IκB proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF-κB to enter the nucleus where it regulates gene expression .

包装清单:

产品编号	产品名称	包装
AF0246	NF-κB p65 Rabbit Polyclonal Antibody	50μl
AZ050	Western一抗稀释液	50ml
-	说明书	1份

保存条件：

NF-κB p65 Rabbit Polyclonal Antibody -20°C保存，Western一抗稀释液-20°C或4°C保存，一年有效。Western一抗稀释液优先推荐4°C保存，长期不使用可以考虑-20°C保存，但冻融可能会导致出现轻微的浑浊和少量不溶物。

注意事项：

- 如果本抗体用于Western blot (WB)、免疫荧光(IF)、免疫细胞化学(ICC)等实验，请注意回收使用过的稀释抗体。回收的抗体通常至少可以重复使用5-10次。稀释后的抗体，包括已经使用过的稀释抗体，请4°C保存。
- 回收后重复使用的抗体，使用方法同新鲜稀释的抗体。如果在重复使用过程中发现抗体出现轻微混浊现象，可以10,000g离心1-3分钟，取上清用于后续检测。如果回收的抗体出现明显的絮状物或长霉长菌等情况，则可以考虑废弃该抗体。
- 提供的Western一抗稀释液也可以用于免疫荧光(IF)、免疫组化(IHC)、免疫细胞化学(ICC)等适当用途。如果希望获得最佳的检测效果，请考虑使用上述检测专用的一抗稀释液。
- 本产品仅限于专业人员的科学研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

使用说明：

请根据抗体的实际用途选择相应的使用方法。

1. Western检测：

- A. 按照推荐的稀释比例用碧云天提供的Western一抗稀释液稀释抗体。
- B. 把经过封闭的蛋白膜与稀释好的一抗4°C缓慢摇动过夜或室温缓慢摇动2小时，确保稀释的抗体至少能在摇动的瞬间覆盖蛋白膜。
- C. 回收稀释的一抗，4°C保存以备下次继续使用。
- D. 按照Western的实验步骤进行后续的洗涤、二抗孵育、洗涤和检测等操作。具体操作可以参考如下网页：
<http://www.beyotime.com/support/western.htm>

2. 免疫染色：

可以使用碧云天生产的免疫染色一抗稀释液(P0103)稀释抗体，使用后注意回收稀释好的一抗，具体操作可以参考如下网页：<http://www.beyotime.com/support/immunol-staining.htm>

3. 其它实验操作请自行参考适当的protocol进行。

4. 代表性图片：




Fig. 1. Western blot analysis of NF-κB p65 on different cell lysates using anti-NF-κB p65 antibody at 1/1000 dilution. Positive control: Lane1: Hela Lane2:A549 Lane3: PC12 Lane 4: Mouse embryonic Lane5: NIH/3T3 stem cell




Fig. 2. Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-NF-κB p65 antibody. Counter stained with hematoxylin.




Fig. 3. Immunohistochemical analysis of paraffin-




Fig. 4. Flow cytometric analysis of Hela cells with NF-κB p65-FITC.

embedded human lung carcinoma tissue using anti- NF-
kB p65 antibody. Counter stained with hematoxylin.

kB p65 antibody at 1/50 dilution (blue) compared with
an unlabelled control (cells without incubation with
primary antibody; red). Goat anti rabbit IgG (FITC)
was used as the secondary antibody.

相关产品：

产品编号	产品名称	包装
P0006	Bradford蛋白浓度测定试剂盒	1000次
P0010	BCA蛋白浓度测定试剂盒(增强型)	500次
P0012	BCA蛋白浓度测定试剂盒	500次
P0012A	SDS-PAGE凝胶配制试剂盒	1盒
P0012AC	SDS-PAGE凝胶快速配制试剂盒	1盒
P0013	Western及IP细胞裂解液	100ml
P0013B	RIPA裂解液(强)	100ml
P0014B	SDS-PAGE电泳液	10×1L
P0015	SDS-PAGE蛋白上样缓冲液(5X)	2ml
P0018	BeyoECL Plus(超敏ECL化学发光试剂盒)	共100ml
P0018A	BeyoECL Star(特超敏ECL化学发光试剂盒)	共100ml
P0020	显影定影试剂盒	各1升
P0021B	Western转膜液	10×1L
P0023A	Western一抗稀释液	100ml
P0023B	Western封闭液	100ml
P0023C6	Western洗涤液(10X)	10×100ml
P0023D	Western二抗稀释液	100ml
P0025	Western一抗二抗去除液	250ml
P0252	QuickBlock™ Western封闭液	100ml
P0256	QuickBlock™ Western一抗稀释液	100ml
P0258	QuickBlock™ Western二抗稀释液	100ml
P0066	预染蛋白质分子量标准	200μl
P0068	彩色预染蛋白质分子量标准	200μl
P0071	BeyoColor™彩色预染蛋白质分子量标准	200μl
P0098	免疫染色固定液	100ml
P0102	免疫染色封闭液	100ml
P0103	免疫染色一抗稀释液	100ml
P0106L	免疫染色洗涤液(10X)	10×100ml
P0108	免疫荧光染色二抗稀释液	100ml
P0110	免疫染色(非荧光)二抗稀释液	100ml
P0260	QuickBlock™免疫染色封闭液	100ml
P0262	QuickBlock™免疫染色一抗稀释液	100ml
P0265	QuickBlock™免疫荧光染色二抗稀释液	100ml
P0267	QuickBlock™免疫组化染色二抗稀释液	100ml
P0126	抗荧光淬灭封片液	5ml
FFP24	PVDF膜(进口分装, 6.6×8.5cm, 0.2μm)	20张/包装
FFP32	PVDF膜(进口原装, 6.6×8.5cm, 0.45μm)	20张/包装
FFP51	转印滤纸(7.5×10cm)	100片/包装
FF057	X-OMAT BT胶片(原柯达, 5×7英寸)	100张/盒
FFC58	压片暗盒(5×7英寸)	1个

使用本产品的文献：

- Lin Z, Chen B, Wu T, Xu X . Highly Tumorigenic Diffuse Large B Cell Lymphoma Cells Are Produced by Coculture with Stromal Cells. ACTA HAEMATOL-BASEL. 2018;139(4):201-216.
- Hui Wang, Zhengxia Wang, Qingbin Tang . Reduced Expression of microRNA-199a-3p Is Associated With Vascular Endothelial Cell Injury Induced by Type 2 Diabetes Mellitus Exp Ther Med. 2018 Oct;16(4):3639-3645.

3. Gao Y, Zhang X, Yin J, Du Q, Tu Y, Shi J, Xu Y . Castanopsis lamontii Water Extract Shows Potential in Suppressing Pathogens, Lipopolysaccharide-Induced Inflammation and Oxidative Stress-Induced Cell Injury. *Molecules*. 2019 Jan 12;24(2). pii: E273.
4. Gao M, Guo Y, Luo X, Du J, Wang Y, Cao C, Wang J, Han W . Design, biological activity and signaling pathway of bovine consensus omega interferon expressed in Pichia pastoris. *Mol Immunol*. 2019 Feb;106:46-52.
5. Yang F, Jin S, Tang Y . Marine Collagen Peptides Promote Cell Proliferation of NIH-3T3 Fibroblasts via NF-κB Signaling Pathway. *Molecules*. 2019 Nov 19;24(22). pii: E4201.
6. Zou Y, Zhou C, Xu H, Yu J, Ye P, Zhang H, Chen S, Zhao J, Le S, Cui J, Jiang L, Wu J, Xia J . Glibenclamide ameliorates transplant-induced arteriosclerosis and inhibits macrophage migration and MCP-1 expression. *Life Sci*. 2020 Jan 15;241:117141.
7. Yanqiang Zou, Cheng Zhou, Heng Xu, Jizhang Yu, Ping Ye, Hao Zhang, Shanshan Chen, Jing Zhao, Sheng Le, Jikai Cui, Lang Jiang, Jie Wu, Jiahong Xia . Glibenclamide ameliorates transplant-induced arteriosclerosis and inhibits macrophage migration and MCP-1 expression *Life Sci*. 2020 Jan 15;241:117141.
8. Li M, Hu FC, Qiao F, Du ZY, Zhang ML . Sodium acetate alleviated high-carbohydrate induced intestinal inflammation by suppressing MAPK and NF-κB signaling pathways in Nile tilapia (*Oreochromis niloticus*). *FISH SHELLFISH IMMUN*. 2020 Mar;98:758-765.
9. Miao Li, Fang-Chao Hu, Fang Qiao, Zhen-Yu Du, Mei-Ling Zhang . Sodium acetate alleviated high-carbohydrate induced intestinal inflammation by suppressing MAPK and NF-κB signaling pathways in Nile tilapia (*Oreochromis niloticus*) *FISH SHELLFISH IMMUN*. 2020 Mar;98:758-765.
10. Huan Gao, Xiangfeng Wang, Xiaoyu Qu, Jinghui Zhai, Lina Tao, Yueming Zhang, Yanqing Song, Wenrui Zhang . Omeprazole attenuates cisplatin-induced kidney injury through suppression of the TLR4/NF-κB/NLRP3 signaling pathway *Toxicology*. 2020 Jul;440:152487.
11. Shibin Feng, Hongyan Ding, Leihong Liu, Chenglu Peng, Yingying Huang, Fuchao Zhong, Wei Li, Tingting Meng, Jinchun Li, Xichun Wang, Yu Li, Jinjie Wu . Astragalus polysaccharide enhances the immune function of RAW264.7 macrophages via the NF-κB p65/MAPK signaling pathway *Exp Ther Med*. 2021 Jan;21(1):20.
12. Junyu Hou, Zhimin Lei, Linlu Cui, Yun Hou, Long Yang, Ru An, Qimeng Wang, Shengda Li, Hongqin Zhang, Lianshuang Zhang . Polystyrene microplastics lead to pyroptosis and apoptosis of ovarian granulosa cells via NLRP3/Caspase-1 signaling pathway in rats *Ecotoxicol Environ Saf*. 2021 Apr 1;212:112012.
13. Yang Xue, Shendong Zhou, Wan Xie, Meijuan Meng, Nana Ma, Hongzhu Zhang, Yan Wang, Guangjun Chang, Xiangzhen Shen . STIM1-Orail Interaction Exacerbates LPS-Induced Inflammation and Endoplasmic Reticulum Stress in Bovine Hepatocytes through Store-Operated Calcium Entry Genes (Basel). 2022 May 13;13(5):874.
14. Yuqi Ma, Peixia Fan, Rui Zhao, Yinghua Zhang, Xianwei Wang, Weigang Cui . Neuregulin-1 regulates the conversion of M1/M2 microglia phenotype via ErbB4-dependent inhibition of the NF-κB pathway *Mol Biol Rep*. 2022 May;49(5):3975-3986.
15. Xing-Mei Shu, Yang Hu, Xin Fang, Jing Wang, Xiao-Yan Qin, Rongfeng Lan . Salidroside alleviates cadmium-induced toxicity in mice by restoring the notch/HES-1 and RIP1-driven inflammatory signaling axis *Inflamm Res*. 2022 Jun;71(5-6):615-626.
16. Zihan Wang, Jiancong Hu, Junxiong Chen, Jingdan Zhang, Weiqian Li, Yu Tian, Huanliang Liu, Xiangling Yang . ICAT promotes colorectal cancer metastasis via binding to JUP and activating the NF-κB signaling pathway *J Clin Lab Anal*. 2022 Oct;36(10):e24678.
17. Haiyue Dai, Yixiao Wang, Zhenying Fan, Yongli Guo, Jiaqi Chen, Ye Meng, Xin Tong, Mingchun Gao, Junwei Wang . Bovine cyclic GMP-AMP synthase recognizes exogenous double-stranded DNA and activates the STING-depended interferon β production pathway *Dev Comp Immunol*. 2023 Feb;139:104567.

Version 2024.03.12