

pARE-luc (报告基因质粒)


| 产品编号 | 产品名称 | 包装 |
|-------------|-------------------|-------|
| D2112-1μg | pARE-luc (报告基因质粒) | 1μg |
| D2112-100μg | pARE-luc (报告基因质粒) | 100μg |

产品简介：

- pARE-luc (报告基因质粒)是碧云天自行研发的用于检测基于顺式作用元件antioxidant response element(ARE, 抗氧化响应元件)的转录活性水平的报告基因质粒。pARE-luc是以碧云天的pGL6质粒为模板, 在其多克隆位点插入了2个ARE结合位点, 可以高灵敏度地检测顺式作用元件ARE的激活水平, 可以用于检测细胞的抗氧化水平或氧化应激状态。
- pGL6质粒模板是用于在哺乳动物细胞中进行萤火虫萤光素酶(firefly luciferase)报告基因检测的新一代质粒。该报告基因质粒比Promega公司的pGL3系列有了全面的改进, 一方面对于luciferase的编码进行了改进, 确保能更好地在哺乳动物细胞中进行表达, 同时整个质粒中所有可以被预测出的可能的转录因子结合位点全部进行了适当的突变处理, 在保持原有功能不变的情况下, 使各种转录因子在质粒上的非特异性结合降到最低。
- 萤光素、萤光素酶、萤火虫萤光素酶和海肾萤光素酶也经常被称为荧光素、荧光素酶、萤火虫荧光素酶和海肾荧光素酶。
- pARE-luc质粒的主要信息如下:

| | |
|--|-----------|
| Base pairs | 5622 |
| ARE response element (ARE RE) | 32-77 |
| luc2 reporter gene | 123-1775 |
| SV40 late poly(A) signal | 1810-2031 |
| SV40 early enhancer/promoter | 2079-2497 |
| Synthetic neomycin phosphotransferase (Neor) coding region | 2522-3316 |
| Synthetic poly(A) signal | 3341-3389 |
| Reporter Vector primer 4 (RVprimer4) binding region | 3456-3475 |
| ColE1-derived plasmid replication origin | 3713 |
| Synthetic Beta-lactamase (Amp ^r) coding region | 4504-5364 |
| Synthetic poly(A) signal/transcriptional pause site | 5469-5622 |
| Reporter Vector primer 3 (RVprimer3) binding region | 5571-5590 |

- pARE-luc质粒的图谱如下:



➤ pARE-luc的多克隆位点及ARE response element的详细图谱如下:

BglI KpnI NheI XhoI ARE response element
 1 GGCCTAACTG GCCGGTACCG CTAGCCTCGA GACTGAGGGT GACTCAGCAA
 CCGGATTGAC CGGCCATGGC GATCGGAGCT CTGACTCCCA CTGAGTCGTT

BglIII HindIII
 51 AATCACTGAG GGTGACTCAG CAAAATCAGA TCTGCAGAAG CTTGGCAATC
 TTAGTGACTC CCACTGAGTC GTTTTAGTCT AGACGTCTTC GAACCGTTAG

➤ pARE-luc中没有的酶切位点(Restriction enzymes that do not cut pARE-luc)包括:

Aat II Afl II Asc I Ase I Bsa I BsaA I BsiW I
 BspM II BssH II Eco72 I EcoR I EcoR V Mlu I Nde I
 Nru I PflM I Pme I Pml I Psp1406 I PspA I Rsr II
 Sac I Sma I SnaB I Spl I Srf I Tth111 I Vsp I
 Xcm I Xma I

➤ pARE-luc中的单酶切位点(Restriction enzymes that cut pARE-luc once)包括:

| | | | | | |
|----------|------------------|------|-----------|-----------------|-----------|
| Sfi I | GGCCN, NNN`NGGCC | 9 | EcoN I | CCTNN`N, NNAGG | 2997 |
| Bgl I | GCCN, NNN`NGGC | 9 | BsiC I | TT`CG, AA | 3392 |
| Acc65 I | G`GTAC, C | 15 | BstB I | TT`CG, AA | 3392 |
| Asp718 | G`GTAC, C | 15 | Sal I | G`TCGA, C | 3406 |
| Kpn I | G, GTAC`C | 19 | Afl III | A`CRYG, T | 3656 |
| Nhe I | G`CTAG, C | 21 | ApaL I | G`TGCA, C | 3970 |
| PaeR7 I | C`TCGA, G | 27 | HgiE II | ACCNNNNNNGGT | -1/134235 |
| Xho I | C`TCGA, G | 27 | Not I | GC`GGCC, GC | 4476 |
| Bgl II | A`GATC, T | 79 | BstX I | CCAN, NNNN`NTGG | 4500 |
| Hind III | A`AGCT, T | 89 | BstE II | G`GTNAC, C | 4503 |
| BsrG I | T`GTAC, A | 614 | Ahd I | GACNN, N`NNGTC | 4578 |
| Dra III | CAC, NNN`GTG | 1270 | Bsu36 I | CC`TNA, GG | 4934 |
| Gsu I | CTGGAG 21/19 | 1503 | Pvu I | CG, AT`CG | 4948 |
| Bpm I | CTGGAG 22/20 | 1504 | Sac II | CC, GC`GG | 4972 |
| Apo I | R`AATT, Y | 1886 | Bst1107 I | GTA TAC | 5088 |
| Mun I | C`AATT, G | 1950 | Xca I | GTA TAC | 5088 |
| BamH I | G`GATC, C | 2043 | Spe I | A`CTAG, T | 5407 |
| Stu I | AGG CCT | 2476 | BsmB I | CGTCTC 7/11 | 5420 |

➤ pARE-luc质粒中推荐使用的测序引物序列如下:

RVprimer3 (5571-5590):
 CTA GCA AAA TAG GCT GTC CC

➤ pARE-luc的全序列信息请参考碧云天的网站上该质粒的信息。

包装清单 :

| 产品编号 | 产品名称 | 包装 |
|-------------|-------------------|-------|
| D2112-1μg | pARE-luc (报告基因质粒) | 1μg |
| D2112-100μg | pARE-luc (报告基因质粒) | 100μg |
| — | 说明书 | 1份 |

保存条件 :

-20°C保存。

注意事项:

- 本质粒未经碧云天书面许可不得用于任何商业用途, 也不得移交给订货人所在实验室外的任何个人或单位。
- 本产品仅限于专业人员的科学研究用, 不得用于临床诊断或治疗, 不得用于食品或药品, 不得存放于普通住宅内。
- 为了您的安全和健康, 请穿实验服并戴一次性手套操作。

使用说明 :

1. 首次使用1μg包装的本产品时请先取少量本质粒转化大肠杆菌, 进行质粒小量、中量或大量抽提后再用于后续用途。抽提获得的质粒可以通过酶切电泳进行鉴定, 或通过测序进行鉴定。
2. 100μg包装的本产品质粒浓度为0.1μg/μl, 共1ml。可以直接用于酶切或者转染细胞。
3. pARE-luc可以用常规的细胞转染方法转染细胞。检测时可以采用碧云天的萤火虫荧光素酶报告基因检测试剂盒(RG005/RG006)或双萤光素酶报告基因检测试剂盒(RG027/RG028)。

使用本产品的文献 :

1. Li C, Pan Z, Xu T, Zhang C, Wu Q, Niu Y. Puerarin induces the upregulation of glutathione levels and nuclear translocation of Nrf2 through

- PI3K/Akt/GSK-3 β signaling events in PC12 cells exposed to lead. *Neurotoxicol Teratol*. 2014 Sep 4;46C:1-9.
2. Huang K, Chen C, Hao J, Huang J, Wang S, Liu P, Huang H. Polydatin promotes Nrf2-ARE anti-oxidative pathway through activating Sirt1 to resist AGEs-induced upregulation of fibronectin and transforming growth factor- β 1 in rat glomerular mesangial cells. *Mol Cell Endocrinol*. 2015 Jan 5;399:178-89.
 3. Zou Y, Wang R, Guo H, Dong M. Phytoestrogen β -Ecdysterone Protects PC12 Cells Against MPP+-Induced Neurotoxicity In Vitro: Involvement of PI3K-Nrf2-Regulated Pathway. *Toxicol Sci*. 2015 Sep;147(1):28-38.
 4. Shan JL, He HT, Li MX, Zhu JW, Cheng Y, Hu N, Wang G, Wang D, Yang XQ, He Y, Xiao HL, Tong WD, Yang ZZ. APE1 promotes antioxidant capacity by regulating Nrf-2 function through a redox-dependent mechanism. *Free Radic Biol Med*. 2015 Jan;78:11-22.
 5. Shi Z, Niu Y, Wang Q, Shi L, Guo H, Liu Y, Zhu Y, Liu S, Liu C, Chen X, Zhang R. Reduction of DNA damage induced by titanium dioxide nanoparticles through Nrf2 in vitro and in vivo. *J Hazard Mater*. 2015 Nov 15;298:310-9.
 6. Wang P, Gao YM, Sun X, Guo N, Li J, Wang W, Yao LP, Fu YJ. Hepatoprotective effect of 2'-O-galloylhyperin against oxidative stress-induced liver damage through induction of Nrf2/ARE-mediated antioxidant pathway. *Food Chem Toxicol*. 2017 Apr;102:129-142.
 7. Lu C, Zou Y, Liu Y, Niu Y. Rosmarinic acid counteracts activation of hepatic stellate cells via inhibiting the ROS-dependent MMP-2 activity: Involvement of Nrf2 antioxidant system. *Toxicol Appl Pharmacol*. 2017 Mar 1;318:69-78.
 8. Dai J, Gu L, Su Y, Wang Q, Zhao Y, Chen X, Deng H, Li W, Wang G, Li K. Inhibition of curcumin on influenza A virus infection and influenzal pneumonia via oxidative stress, TLR2/4, p38/JNK MAPK and NF- κ B pathways. *Int Immunopharmacol*. 2018 Jan;54:177-187.
 9. Zhang H, Zheng W, Feng X, Yang F, Qin H, Wu S, Hou DX, Chen J. Nrf2-ARE Signaling Acts as Master Pathway for the Cellular Antioxidant Activity of Fisetin. *Molecules*. 2019 Feb 15;24(4). pii: E708
 10. Xiang Y, Fan X, Zhao M, Guo Q, Guo S. CKIP-1 alleviates oxygen-glucose deprivation/reoxygenation-induced apoptosis and oxidative stress in cultured hippocampal neurons by downregulating Keap1 and activating Nrf2/ARE signaling. *Eur J Pharmacol*. 2019 Apr 5;848:140-149
 11. Ma B, Zhang J, Zhu Z, Zhao A, Zhou Y, Ying H, Zhang Q. Luteolin Ameliorates Testis Injury and Blood-Testis Barrier Disruption through the Nrf2 Signaling Pathway and by Upregulating Cx43. *Mol Nutr Food Res*. 2019 May;63(10):e1800843
 12. Ren J, Li J, Hu J, Yu H, Yang Q, Mu J. Overexpression of CKIP-1 alleviates hypoxia-induced cardiomyocyte injury by up-regulating Nrf2 antioxidant signaling via Keap1 inhibition. *Biochimie*. 2019 Aug;163:163-170
 13. Tian X, Cong F, Guo H, Fan J, Chao G, Song T. Downregulation of Bach1 protects osteoblasts against hydrogen peroxide-induced oxidative damage in vitro by enhancing the activation of Nrf2/ARE signaling. *CHEM-BIOL INTERACT*. 2019 Aug 25;309:108706
 14. Zhang Z, Wang N, Zhang Y, Zhao J, Lv J. Downregulation of microRNA-302b-3p relieves oxygen-glucose deprivation/re-oxygenation induced injury in murine hippocampal neurons through up-regulating Nrf2 signaling by targeting fibroblast growth factor. *CHEM-BIOL INTERACT*. 2019 Aug 25;309:108705
 15. Yang Y, Tian T, Wang Y, Li Z, Xing K, Tian G. SIRT6 protects vascular endothelial cells from angiotensin II-induced apoptosis and oxidative stress by promoting the activation of Nrf2/ARE signaling. *Eur J Pharmacol*. 2019 Sep 15;859:172516
 16. Zuo H, Wang S, Feng J, Liu X. BRD4 contributes to high-glucose-induced podocyte injury by modulating Keap1/Nrf2/ARE signaling. *Biochimie*. 2019 Oct;165:100-107
 17. Tianrong Xun, Zhufen Lin, Xia Zhan, Shaolian Song, Liqian Mo, Haixing Feng, Qin Yang, Dan Guo, Xixiao Yang. Advanced oxidation protein products upregulate efflux transporter expression and activity through activation of the Nrf-2-mediated signaling pathway in vitro and in vivo. *Eur J Pharm Sci*. 2020 Apr 18;149:105342.
 18. Yin Du, Xiaozhen Ma, Lei Ma, Siyuan Li, Juan Zheng, Junlin Lv, Long Cui, Jianrui Lv. Inhibition of microRNA-148b-3p alleviates oxygen-glucose deprivation/reoxygenation-induced apoptosis and oxidative stress in HT22 hippocampal neuron via reinforcing Sestrin2/Nrf2 signalling. *CLIN EXP PHARMACOL P*. 2020 Apr;47(4):561-570.
 19. Xiaoqing Yan, Yue Su, Xia Fan, Hui Chen, Zixian Lu, Zijuan Liu, Yingjian Li, Mei Yi, Guigui Zhang, Chunjie Gu, Kai Wang, Jiamin Wu, Da Sun, Yikai Zhang, Chi Zhang, Xiaozhen Dai, Chao Zheng. Liraglutide Improves the Angiogenic Capability of EPC and Promotes Ischemic Angiogenesis in Mice under Diabetic Conditions through an Nrf2-Dependent Mechanism. *Cells*. 2022 Nov 29;11(23):3821.

Version 2024.03.12