



碧云天生物技术/Beyotime Biotechnology  
 订货热线: 400-1683301 或 800-8283301  
 订货 e-mail: order@beyotime.com  
 技术咨询: info@beyotime.com  
 网址: http://www.beyotime.com

## Recombinant Human IL-13, Variant

产品编号	产品名称	包装
P5183-10μg	Recombinant Human IL-13, Variant	10μg
P5183-100μg	Recombinant Human IL-13, Variant	100μg
P5183-500μg	Recombinant Human IL-13, Variant	500μg

### 产品简介:

Species	Gene ID	Accession	Source	Length	MW	Tag
Human	3596	P35225	<i>E. coli</i>	112aa	12.3kDa	—

About this protein	
Name	Recombinant Human IL-13, Variant (Recombinant Human Interleukin-13, Variant; rHuIL-13, Variant); 重组人白细胞介素13变体
Synonyms	IL-13; P600
Purity	>97% by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 0.5ng/ml, corresponding to a specific activity of >2.0×10 <sup>6</sup> IU/mg.
Physical Appearance	Sterile Filtered lyophilized (freeze-dried) liquid-like powder
Formulation	Lyophilized from a 0.2μm filtered concentrated solution in PBS, pH7.2, with 5% trehalose.
Endotoxin	Less than 1EU/μg of rHuIL-13, Variant as determined by LAL method.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions.
Category	Cytokine
Background	Human Interleukin-13 (IL-13) is expressed by the IL13 gene located on the chromosome 5. IL-13 is a 132 amino acid protein containing a proposed 20 amino acid signal peptide, and shares approximately 30% amino acid sequence homology to human IL-4. Additionally, the two cytokines exhibit overlapping biological activities. Human IL-13 is produced by activated Th0, Th1-like Th2-like and CD8 T cells. Similarly to IL-4, IL-13 has multiple effects on the differentiation and functions of monocytes/macrophages. IL-13 can suppress the cytotoxic functions of monocytes/macrophages. IL-13 variant is a natural variant of IL-13 and has a Q instead of R112 of wild type, which associated with increased risk for asthma development and at homozygosity associated with higher levels of serum total IgE in some allergic rhinitis patients.
Amino Acid Sequence	GPVPPSTALR ELIEELVNIT QNQKAPLCNG SMVWSINLTA GMYCAALES L INVSGCSAIE KTQRMLSGFC PHKVSAGQFS SLHVRDTKIE VAQFVKDLLL HLKKLFRREGQ FN

### 包装清单:

产品编号	产品名称	包装
P5183-10μg	Recombinant Human IL-13, Variant	10μg
P5183-100μg	Recombinant Human IL-13, Variant	100μg
P5183-500μg	Recombinant Human IL-13, Variant	500μg
—	说明书	1份

### 保存条件:

-20°C或更低温度保存，至少一年有效。由于蛋白的每次冻融均会引起部分失活，所以首次配制相应浓度的储存液后(请根据产品简介中Reconstitution一栏的信息配制储存液)，须分装后-20°C或更低温度冻存，以避免反复冻融。

### 注意事项：

- 由于有些塑料管壁对某些蛋白有较强的吸附作用，溶液中的蛋白很容易粘附在管壁上，并且粘附后的蛋白很难与管壁分离。而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点，使细胞因子或重组蛋白不会粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- 本产品在冻干时添加了海藻糖作为保护剂，此时冻干后本产品看起来仍然像是液体，属于正常现象。在冻干工艺上，海藻糖作为常用的冻干保护剂，可显著阻止蛋白质二级结构的改变，防止冻干过程中的蛋白变性。
- 本产品仅限于专业人员的科学研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

### 使用说明：

1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明，碧云天相关产品均为冻干粉，由于微量的蛋白在冻干过程中沉积在管内，形成很薄或不可见的蛋白层，所以在打开管盖前，我们建议在离心机中约8,000-12,000g离心10-30秒，使附着在管盖或管壁上的蛋白聚集于管底。
2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是非常容易溶解的，一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子，即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡，以免蛋白变性而失活。
3. 具体的最佳工作浓度请自行参考相关文献，或者根据实验目的，以及特定细胞和动物，通过实验进行摸索和优化。

Version 2024.01.01