



碧云天生物技术/Beyotime Biotechnology
订货热线：400-168-3301或800-8283301
订货e-mail：order@beyotime.com
技术咨询：info@beyotime.com
网址：http://www.beyotime.com

Recombinant Murine FGF-21

| 产品编号 | 产品名称 | 包装 |
|-------------|---------------------------|-------|
| P6101-5μg | Recombinant Murine FGF-21 | 5μg |
| P6101-25μg | Recombinant Murine FGF-21 | 25μg |
| P6101-100μg | Recombinant Murine FGF-21 | 100μg |
| P6101-1mg | Recombinant Murine FGF-21 | 1mg |

产品简介：

| Species | Gene ID | Accession | Source | Length | MW | Tag |
|---------|---------|-----------|----------------|--------|---------|-----|
| Murine | 56636 | Q9JJN1 | <i>E. coli</i> | 182aa | 19.9kDa | — |

| About this protein | |
|---------------------|--|
| Name | Recombinant Murine FGF-21 (Recombinant Murine Fibroblast Growth Factor-21; rMuFGF-21); 重组小鼠成纤维细胞生长因子21 |
| Synonyms | FGF-21; fibroblast growth factor 21 |
| Purity | >97% by SDS-PAGE and HPLC analyses. |
| Biological Activity | Fully biologically active when compared to standard. The ED50 as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 0.5μg/ml, corresponding to a specific activity of >2.0×10 ³ IU/mg in the presence of 5μg/ml of rMuKlotho-β and 10μg/ml of heparin. |
| Physical Appearance | Sterile Filtered White lyophilized (freeze-dried) powder. |
| Formulation | Lyophilized from a 0.2μm filtered concentrated solution in 3X PBS, pH7.4. |
| Endotoxin | Less than 1EU/μg of rMuFGF-21 as determined by LAL method. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions. |
| Category | Cytokine |
| Background | Fibroblast growth factor-21 (FGF-21) belongs to the large FGF family and it is specifically induced by HMGCS2 activity. In mice, brown adipose tissue becomes a source of systemic FGF21 after cold exposure. FGF-21 stimulates glucose uptake in differentiated adipocytes via the induction of glucose transporter SLC2A1/GLUT1 expression (but not SLC2A4/GLUT4 expression) and the activity depends on the presence of KLB. FGF-21, in the presence of betaKlotho as a protein cofactor, signals through the FGFR 1c and 4 receptors. Recombinant murine FGF-21 contains 182 amino acids residues and show limited binding to heparin. In addition, Murine FGF-21 respectively shows 81% and 92% a.a. identity to human and rat FGF-21, and it show activity on human and rat cells. |
| Amino Acid Sequence | AYPIPDSSPL LQFGGQVRQR YLYTDDQDT EAHLEIREDG TVVGAAHRSP ESLLELKALK PGVIQILGVK ASRFLCQQPD GALYGSPHFD PEACSFRELL LEDGYNVYQS EAHGLPLRLP QKDSPNQDAT SWGPVRFLPM PGLLHEPQDQ AGFLPPEPPD VGSSDPLSMV EPLQGRSPSY AS |

包装清单：

| 产品编号 | 产品名称 | 包装 |
|-------------|---------------------------|-------|
| P6101-5μg | Recombinant Murine FGF-21 | 5μg |
| P6101-25μg | Recombinant Murine FGF-21 | 25μg |
| P6101-100μg | Recombinant Murine FGF-21 | 100μg |
| P6101-1mg | Recombinant Murine FGF-21 | 1mg |
| — | 说明书 | 1份 |

保存条件：

-20°C或更低温度保存，至少一年有效。由于蛋白的每次冻融均会引起部分失活，所以首次配制成相应浓度的储存液后(请根据产品简介中Reconstitution一栏的信息配制储存液)，须分装后-20°C或更低温度冻存，以避免反复冻融。

注意事项：

- 由于有些塑料管壁对某些蛋白有较强的吸附作用，溶液中的蛋白很容易粘附在管壁上，并且粘附后的蛋白很难与管壁分离。而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点，使细胞因子或重组蛋白不会粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- 本产品仅限于专业人员的科学的研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

使用说明：

1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明，碧云天相关产品均为冻干粉，由于微量的蛋白在冻干过程中沉积在管内，形成很薄或不可见的蛋白层，所以在打开管盖前，我们建议在离心机中约8,000-12,000g离心10-30秒，使附着在管盖或管壁上的蛋白聚集于管底。
2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是非常容易溶解的，一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子，即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡，以免蛋白变性而失活。
3. 具体的最佳工作浓度请自行参考相关文献，或者根据实验目的，以及特定细胞和动物，通过实验进行摸索和优化。

Version 2017.02.10