



碧云天生物技术/Beyotime Biotechnology  
 订货热线: 400-1683301 或 800-8283301  
 订货 e-mail: order@beyotime.com  
 技术咨询: info@beyotime.com  
 网址: http://www.beyotime.com

## Recombinant Human GCP-2/CXCL6

产品编号	产品名称	包装
P6515-5μg	Recombinant Human GCP-2/CXCL6	5μg
P6515-100μg	Recombinant Human GCP-2/CXCL6	100μg
P6515-500μg	Recombinant Human GCP-2/CXCL6	500μg

### 产品简介:

Species	Gene ID	Accession	Source	Length	MW	Tag
Human	6372	P80162	<i>E. coli</i>	72aa	7.9kDa	—

About this protein	
Name	Recombinant Human GCP-2/CXCL6 (Recombinant Human Granulocyte Chemotactic Protein 2/CXCL6; rHuGCP-2/CXCL6); 重组人粒细胞趋化蛋白2/CXCL6
Synonyms	GCP-2; chemokine (C-X-C motif) ligand 6 (granulocyte chemotactic protein 2); Chemokine alpha 3; CKA-3Granulocyte chemotactic protein 2; GCP2member 6 (granulocytechemotactic protein 2); GCP-2Small-inducible cytokine B6; Small inducible cytokine subfamily B (Cys-X-Cys), member b
Purity	>98% by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 10-50ng/ml.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation	Lyophilized from a 0.2μm filtered concentrated solution in PBS, pH7.4.
Endotoxin	Less than 0.1EU/μg of rHuGCP-2/CXCL6 as determined by LAL method.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions.
Category	Chemokines
Background	GCP-2 (granulocyte chemotactic protein-2) is a CXC chemokine. Among human CXC chemokines, GCP-2 is most closely related to ENA-78. The structure and sequence of the genes for human GCP-2 and ENA-78 also exhibit close similarity, suggesting the two genes may have originated from a recent gene duplication. LIX (LPS-induced CXC chemokine) was initially cloned as a gene induced by LPS in mouse fibroblasts. The mouse protein designated GCP-2, because of its amino acid sequence similarity (60%) to human GCP-2, is identical to the LIX protein sequence.
Amino Acid Sequence	VLTELRTCL RVTLRVNPKT IGKLQVFPAG PQCSKVEVVA SLKNGKQVCL DPEAPFLKKV IQKILDSGNK KN

### 包装清单:

产品编号	产品名称	包装
P6515-5μg	Recombinant Human GCP-2/CXCL6	5μg
P6515-100μg	Recombinant Human GCP-2/CXCL6	100μg
P6515-500μg	Recombinant Human GCP-2/CXCL6	500μg
—	说明书	1份

### 保存条件:

-20°C或更低温度保存, 至少一年有效。由于蛋白的每次冻融均会引起部分失活, 所以首次配制相应浓度的储存液后(请根据产品简介中Reconstitution一栏的信息配制储存液), 须分装后-20°C或更低温度冻存, 以避免反复冻融。

### 注意事项:

- 由于有些塑料管壁对某些蛋白有较强的吸附作用，溶液中的蛋白很容易粘附在管壁上，并且粘附后的蛋白很难与管壁分离。而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点，使细胞因子或重组蛋白不会粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- 本产品仅限于专业人员的科学研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

#### 使用说明：

1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明，碧云天相关产品均为冻干粉，由于微量的蛋白在冻干过程中沉积在管内，形成很薄或不可见的蛋白层，所以在打开管盖前，我们建议在离心机中约8,000-12,000g离心10-30秒，使附着在管盖或管壁上的蛋白聚集于管底。
2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是很容易溶解的，一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子，即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡，以免蛋白变性而失活。
3. 具体的最佳工作浓度请自行参考相关文献，或者根据实验目的，以及特定细胞和动物，通过实验进行摸索和优化。

Version 2023.12.20