



碧云天生物技术/Beyotime Biotechnology
订货热线: 400-1683301 或 800-8283301
订货 e-mail: order@beyotime.com
技术咨询: info@beyotime.com
网址: http://www.beyotime.com

Recombinant Human MDC/CCL22

| 产品编号 | 产品名称 | 包装 |
|-------------|-----------------------------|-------|
| P6659-5μg | Recombinant Human MDC/CCL22 | 5μg |
| P6659-100μg | Recombinant Human MDC/CCL22 | 100μg |
| P6659-500μg | Recombinant Human MDC/CCL22 | 500μg |

产品简介:

| Species | Gene ID | Accession | Source | Length | MW | Tag |
|---------|---------|-----------|----------------|--------|--------|-----|
| Human | 6367 | O00626 | <i>E. coli</i> | 69aa | 8.1kDa | — |

| About this protein | |
|---------------------|---|
| Name | Recombinant Human MDC/CCL22 (Recombinant Human Macrophage-Derived Chemokine/CCL22; rHuMDC/CCL22); 重组人巨噬细胞来源趋化因子/CCL22 |
| Synonyms | ABCD-1; MDC; STCP-1; A-152E5.1; ABCD-1; CC chemokine STCP-1; C-C motif chemokine 22; chemokine (C-C motif) ligand 22; DC/B-CK; Macrophage-derived chemokine; MDCStimulated T-cell chemotactic protein 1; MGC34554; SCYA22MDC(1-69); small inducible cytokine A22; small inducible cytokine subfamily A (Cys-Cys), member 22; Small-inducible cytokine A22; STCP-1; stimulated T cell chemotactic protein 1 |
| Purity | >97% by SDS-PAGE and HPLC analyses. |
| Biological Activity | Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 10-100ng/ml. |
| Physical Appearance | Sterile Filtered White lyophilized (freeze-dried) powder. |
| Formulation | Lyophilized from a 0.2μm filtered concentrated solution in 20mM PB, pH7.4, 500mM NaCl. |
| Endotoxin | Less than 1EU/μg of rHuMDC/CCL22 as determined by LAL method. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions. |
| Category | Chemokines |
| Background | CCL22 is a protein that in humans is encoded by the CCL22 gene, which locates on the Chr. 16. The protein is highly expressed in macrophage, monocyte-derived dendritic cell and thymus, additionally, also detected in the tissues of thymus, lymph node and appendix. CCL22 can bind to CCR4, and is a chemoattractant for monocytes, monocyte-derived dendritic cells, and natural killer cells, but not for neutrophils, eosinophils, and resting T-lymphocytes. After secreted from monocyte-derived dendritic cells, the protein can be proteolytic cleaved into three forms: MDC (3-69), MDC (5-69), MDC (7-69). |
| Amino Acid Sequence | GPYGANMEDS VCCRDYVRYR LPLRVVKHFY WTSDSCPRPG VVLLTFRDKE ICADPRVPWV KMILNKLSQ |

包装清单:

| 产品编号 | 产品名称 | 包装 |
|-------------|-----------------------------|-------|
| P6659-5μg | Recombinant Human MDC/CCL22 | 5μg |
| P6659-100μg | Recombinant Human MDC/CCL22 | 100μg |
| P6659-500μg | Recombinant Human MDC/CCL22 | 500μg |
| — | 说明书 | 1份 |

保存条件:

-20°C或更低温度保存，至少一年有效。由于蛋白的每次冻融均会引起部分失活，所以首次配制相应浓度的储存液后(请根据产品简介中Reconstitution一栏的信息配制储存液)，须分装后-20°C或更低温度冻存，以避免反复冻融。

注意事项：

- 由于有些塑料管壁对某些蛋白有较强的吸附作用，溶液中的蛋白很容易粘附在管壁上，并且粘附后的蛋白很难与管壁分离。而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点，使细胞因子或重组蛋白不会粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- 本产品仅限于专业人员的科学的研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

使用说明：

1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明，碧云天相关产品均为冻干粉，由于微量的蛋白在冻干过程中沉积在管内，形成很薄或不可见的蛋白层，所以在打开管盖前，我们建议在离心机中约8,000-12,000g离心10-30秒，使附着在管盖或管壁上的蛋白聚集于管底。
2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是非常容易溶解的，一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子，即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡，以免蛋白变性而失活。
3. 具体的最佳工作浓度请自行参考相关文献，或者根据实验目的，以及特定细胞和动物，通过实验进行摸索和优化。

Version 2024.01.03