



碧云天生物技术/Beyotime Biotechnology  
 订货热线: 400-1683301 或 800-8283301  
 订货 e-mail: order@beyotime.com  
 技术咨询: info@beyotime.com  
 网址: http://www.beyotime.com

## Recombinant Murine MCP-2/CCL8

| 产品编号        | 产品名称                          | 包装    |
|-------------|-------------------------------|-------|
| P6794-5μg   | Recombinant Murine MCP-2/CCL8 | 5μg   |
| P6794-100μg | Recombinant Murine MCP-2/CCL8 | 100μg |
| P6794-500μg | Recombinant Murine MCP-2/CCL8 | 500μg |

### 产品简介:

| Species | Gene ID | Accession | Source         | Length | MW     | Tag |
|---------|---------|-----------|----------------|--------|--------|-----|
| Murine  | 20307   | Q9Z121    | <i>E. coli</i> | 74aa   | 8.5kDa | —   |

| About this protein  |   |
|---------------------|---|
| Name                | Recombinant Murine MCP-2/CCL8 (Recombinant Murine Monocyte Chemotactic Protein-2/CCL8; rMuMCP-2/CCL8); 重组小鼠单核细胞趋化蛋白2/CCL8   |
| Synonyms            | MCP-2; C-C motif chemokine 8; chemokine (C-C motif) ligand 8; HC14; MCP2; MCP-2; member 8 (monocyte chemotactic protein 2); monocyte chemoattractant protein 2; SCYA10; small-inducible cytokine A8   |
| Purity              | >97% by SDS-PAGE and HPLC analyses.   |
| Biological Activity | Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood monocytes is in a concentration range of 10-100ng/ml.   |
| Physical Appearance | Sterile Filtered White lyophilized (freeze-dried) powder.   |
| Formulation         | Lyophilized from a 0.2μm filtered concentrated solution in 20mM PB, pH7.4, 150mM NaCl.  |
| Endotoxin           | Less than 1EU/μg of rMuMCP-2/CCL8 as determined by LAL method.  |
| Reconstitution      | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions.  |
| Category            | Chemokines  |
| Background          | Murine CCL8, also known as monocyte chemotactic protein 2 (MCP-2), is belonging to the CC chemokine family and is encoded by the gene CCL8 in mouse. MCP-2 has two homogeneous MCP-1 (CCL2) and MCP-3 (CCL7). These three MCPs were found by IL-1-beta triggered human MG-63 osteosarcoma cells. CCL8 shares 62% amino acid sequence identity with MCP-1, and shares 58% amino acid sequence identity with MCP-2. CCL8 has chemotactic function for monocytes, eosinophils and neutrophils. In addition, it can also chemoattract activated NK cells. |
| Amino Acid Sequence | GPDKAPVTCC FHVVLKLIPL RVLKSYERIN NIQCPMEAVV FQTKQGMSLC VDPTQKWVSE YMEILDQKSQ ILQP   |

### 包装清单:

| 产品编号        | 产品名称                          | 包装    |
|-------------|-------------------------------|-------|
| P6794-5μg   | Recombinant Murine MCP-2/CCL8 | 5μg   |
| P6794-100μg | Recombinant Murine MCP-2/CCL8 | 100μg |
| P6794-500μg | Recombinant Murine MCP-2/CCL8 | 500μg |
| —           | 说明书                           | 1份    |

### 保存条件:

-20°C或更低温度保存, 至少一年有效。由于蛋白的每次冻融均会引起部分失活, 所以首次配制相应浓度的储存液后(请根据产品简介中Reconstitution一栏的信息配制储存液), 须分装后-20°C或更低温度冻存, 以避免反复冻融。

## 注意事项：

- 由于有些塑料管壁对某些蛋白有较强的吸附作用，溶液中的蛋白很容易粘附在管壁上，并且粘附后的蛋白很难与管壁分离。而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点，使细胞因子或重组蛋白不会粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- 本产品仅限于专业人员的科学研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

## 使用说明：

1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明，碧云天相关产品均为冻干粉，由于微量的蛋白在冻干过程中沉积在管内，形成很薄或不可见的蛋白层，所以在打开管盖前，我们建议在离心机中约8,000-12,000g离心10-30秒，使附着在管盖或管壁上的蛋白聚集于管底。
2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是非常容易溶解的，一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子，即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡，以免蛋白变性而失活。
3. 具体的最佳工作浓度请自行参考相关文献，或者根据实验目的，以及特定细胞和动物，通过实验进行摸索和优化。

Version 2023.12.14