





碧云天生物技术/Beyotime Biotechnology 订货热线: 400-168-3301或800-8283301

订货e-mail: order@beyotime.com 技术咨询: info@beyotime.com 网址: http://www.beyotime.com

Recombinant Human SHH

| 产品编号 | 产品名称 | 包装 |
|-------------|-----------------------|-------|
| P7212-5µg | Recombinant Human SHH | 5μg |
| P7212-25μg | Recombinant Human SHH | 25μg |
| P7212-100μg | Recombinant Human SHH | 100μg |
| P7212-1mg | Recombinant Human SHH | 1mg |

产品简介:

| Species | Gene ID | Accession | Source | Length | MW | Tag |
|---------|---------|-----------|---------|--------|---------|-----|
| Human | 6469 | Q15465 | E. coli | 176aa | 19.8kDa | _ |

| About this protein | 1 |
|------------------------|--|
| Name | Recombinant Human SHH (Recombinant Human Sonic Hedgehog Protein N-Terminus; rHuSHH); 重组人音 渭因子 |
| Synonyms | Shh; HHG1; HHG-1; HLP3; HPE3; MCOPCB5sonic hedgehog (Drosophila) homolog; SMMCIsonic hedgehog homolog (Drosophila); sonic hedgehog; sonic hedgehog homolog; sonic hedgehog protein; TPT; TPTPS |
| Purity | >98% by SDS-PAGE and HPLC analyses. |
| Biological Activity | Fully biologically active when compared to standard. The ED50 as determined by inducing alkaline phosphatase production of murine C3H/10T1/2 cells is less than $1\mu g/ml$, corresponding to a specific activity of $>1.0\times10^3 IU/mg$. |
| Physical Appearance | Sterile Filtered White lyophilized (freeze-dried) powder. |
| Formulation | Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH7.4, 150mM NaCl. |
| Endotoxin | Less than 1EU/µg of rHuSHH as determined by LAL method. |
| Reconstitution | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤-20°C. Further dilutions should be made in appropriate buffered solutions. |
| Category | Others |
| Background | Sonic Hedgehog (SHH) is one of three proteins of the Hedgehog (Hh) family, which also contains Desert Hedgehog (DHH) and Indian Hedgehog (IHH). The three members share a high degree of amino-acid sequence identity (e.g., SHH and IHH are 93% identical). SHH is expressed in fetal intestine, liver, lung, and kidney, but not in adult tissues. The protein consists of 462a.a. with a 23a.a. signal peptide at N-terminus, and is further cleaved into SHH N-Terminus and C-Terminus. SHH has the most critical roles in development, acting as a morphogen involved in patterning many systems, including the limb and midline structures in the brain, spinal cord, the thalamus by the zona limitans intrathalamica and the teeth. In the absence of Sonic HedgeHog, patched receptor represses the constitutive signaling activity of smoothened. SHH-N retains all known signaling capabilities, and can be lipid-modified without receptor affinity reducing, but has more potent than the unmodified form. The rHuSHH has an N-terminal Ile-Val-Ile sequence substituted for the natural occurring chemically modified Cys residue. |
| Amino Acid Sequence | IVIGPGRGFG KRRHPKKLTP LAYKQFIPNV AEKTLGASGR YEGKISRNSE RFKELTPNYN PDIIFKDEEN TGADRLMTQR CKDKLNALAI SVMNQWPGVK LRVTEGWDED GHHSEESLHY EGRAVDITTS DRDRSKYGML ARLAVEAGFD WVYYESKAHI HCSVKAENSV AAKSGG |

包装清单:

| 产品编号 | 产品名称 | 包装 |
|------------|-----------------------|------|
| P7212-5μg | Recombinant Human SHH | 5μg |
| P7212-25μg | Recombinant Human SHH | 25μg |

| Ρ7212-100μg | Recombinant Human SHH | 100μg |
|-------------|-----------------------|-------|
| P7212-1mg | Recombinant Human SHH | 1mg |
| _ | 说明书 | 1份 |

保存条件:

-20℃或更低温度保存,至少一年有效。由于蛋白的每次冻融均会引起部分失活,所以首次配制成相应浓度的储存液后(请根 据产品简介中Reconstitution一栏的信息配制储存液),须分装后-20℃或更低温度冻存,以避免反复冻融。

注意事项:

- ▶ 由于有些塑料管壁对某些蛋白有较强的吸附作用,溶液中的蛋白很容易粘附在管壁上,并且粘附后的蛋白很难与管壁分离。 而载体蛋白(Carrier protein, 如0.1% BSA等)的主要作用是预先封闭塑料管壁上的蛋白结合位点, 使细胞因子或重组蛋白不会 粘附于管壁。所以一定要使用产品简介中Reconstitution一栏的信息配制储存液。
- ▶ 本产品仅限于专业人员的科学研究用,不得用于临床诊断或治疗,不得用于食品或药品,不得存放于普通住宅内。
- ▶ 为了您的安全和健康,请穿实验服并戴一次性手套操作。

使用说明:

- 1. 收到产品后请立即按照说明书推荐的条件保存。除非特别注明,碧云天相关产品均为冻干粉,由于微量的蛋白在冻干过程 中沉积在管内,形成很薄或不可见的蛋白层,所以在打开管盖前,我们建议在离心机中约8,000-12,000g离心10-30秒,使附 着在管盖或管壁上的蛋白聚集于管底。
- 2. 请根据实验目的并按照产品简介中Reconstitution一栏中的信息配制储存液。大多数细胞因子或重组蛋白的冻干粉是非常容易 溶解的,一般用移液枪的枪头轻吹几下或者轻轻摇晃瓶子,即可使细胞因子或重组蛋白完全溶解。请勿用vortex剧烈振荡, 以免蛋白变性而失活。
- 3. 具体的最佳工作浓度请自行参考相关文献,或者根据实验目的,以及特定细胞和动物,通过实验进行摸索和优化。

Version 2017.02.10