



碧云天网站



微信公众号

碧云天生物技术/Beyotime Biotechnology

订货热线: 400-168-3301或800-8283301

订货e-mail: order@beyotime.com

技术咨询: info@beyotime.com

网址: http://www.beyotime.com

Forskolin (腺苷酸环化酶激活剂)

产品编号	产品名称	包装
S1612-1mg	Forskolin (腺苷酸环化酶激活剂)	5mg/ml×0.2ml
S1612-5mg	Forskolin (腺苷酸环化酶激活剂)	5mg
S1612-25mg	Forskolin (腺苷酸环化酶激活剂)	25mg

产品简介:

- Forskolin是一种常用的腺苷酸环化酶激活剂。Forskolin可以通透细胞，激活腺苷酸环化酶(adenylyl cyclase)，从而升高细胞内的cAMP水平。
- Forskolin分子量为410.50，分子式为C₂₂H₃₄O₇，CAS Number: 66575-29-9。本产品纯度大于99%。
- 本Forskolin为进口分装，其中5mg/ml包装产品用DMSO配制，共0.2ml。5mg和25mg包装为粉末装。

包装清单:

产品编号	产品名称	包装
S1612-1mg	Forskolin (腺苷酸环化酶激活剂)	5mg/ml×0.2ml
S1612-5mg	Forskolin (腺苷酸环化酶激活剂)	5mg
S1612-25mg	Forskolin (腺苷酸环化酶激活剂)	25mg
—	说明书	1份

保存条件:

-20°C避光保存。

注意事项:

- 本产品对人体有害，操作时请小心，并注意有效防护以避免直接接触人体或吸入体内。
- 本Forskolin在4°C、冰浴等较低温度情况下会凝固而粘在离心管管底、管壁或管盖内，可以20-25°C水浴温育片刻至全部融解后使用。
- 本产品仅限于专业人员的科学研究用，不得用于临床诊断或治疗，不得用于食品或药品，不得存放于普通住宅内。
- 为了您的安全和健康，请穿实验服并戴一次性手套操作。

使用说明:

1. Forskolin常见使用浓度范围为1-100μM。具体的最佳工作浓度请参考相关文献，或根据实验目的，以及所培养的特定细胞和组织，通过实验进行摸索和优化。

使用本产品的文献:

1. Li Y, He J, Sui S, Hu X, Zhao Y, Li N. Clenbuterol upregulates histone demethylase JHDM2a via the β(2)-adrenoceptor/cAMP/PKA/p-CREB signaling pathway. *Cell Signal*. 2012 Dec;24(12):2297-306.
2. Du B, Luo W, Li R, Tan B, Han H, Lu X, Li D, Qian M, Zhang D, Zhao Y, Liu M. Lgr4/Gpr48 Negatively Regulates TLR2/4-associated Pattern Recognition and Innate Immunity by Targeting CD14Expression. *J Biol Chem*. 2013 May 24;288(21):15131-41.
3. Mu X, He K, Sun H, Zhou X, Chang L, Li X, Chu W, Qiao G, Lu Y. Hydrogen peroxide induces overexpression of angiotensin-converting enzyme in human umbilical vein endothelial cells. *Free Radic Res*. 2013 Feb;47(2):116-22.
4. Peng Y, Shi J, Du X, Wang L, Klocker H, Mo L, Mo Z, Zhang J. Prostaglandin E2 induces stromal cell-derived factor-1 expression in prostate stromal cells by activating protein kinase A and transcription factor Sp1. *Int J Biochem Cell Biol*. 2013 Mar;45(3):521-30.
5. Zheng YF, Liu CF, Lai WF, Xiang Q, Li ZF, Wang H, Lin N. The laxative effect of emodin is attributable to increased aquaporin 3 expression in the colon of mice and HT-29 cells. *Fitoterapia*. 2014 Jul;96:25-32.
6. Hu QX, Dong JH, Du HB, Zhang DL, Ren HZ, Ma ML, Cai Y, Zhao TC, Yin XL, Yu X, Xue T, Xu ZG, Sun JP. Constitutive Gai coupling activity of very large G protein-coupled receptor 1 (VLGR1) and its regulation by PDZD7 protein. *J Biol Chem*. 2014 Aug 29;289(35):24215-25.
7. Liu F, You X, Wang Y, Liu Q, Liu Y, Zhang S, Chen L, Zhang X, Ye L. The oncoprotein HBXIP enhances angiogenesis and growth of breast cancer through modulating FGF8 and VEGF. *Carcinogenesis*. 2014 May;35(5):1144-53.
8. Lv P, Wang Y, Ma J, Wang Z, Li JL, Hong CS, Zhuang Z, Zeng YX. Inhibition of protein phosphatase 2A with a small molecule LB100 radiosensitizes nasopharyngeal carcinoma xenografts by inducing mitotic catastrophe and blocking DNA damage repair. *Oncotarget*. 2014 Sep 15;5(17):7512-24.
9. Sun F, Ding XP, An SM, Tang YB, Yang XJ, Teng L, Zhang C, Shen Y, Chen HZ, Zhu L. Adrenergic DNA damage of embryonic pluripotent cells via β2 receptor signalling. *Sci Rep*. 2015 Oct 30;5:15950.
10. Sun F, Yang XJ, Lv HY, Tang YB, An SM, Ding XP, Li WB, Teng L, Shen Y, Chen HZ, Zhu L. β2-Adrenoreceptor-Mediated Proliferation Inhibition of Embryonic Pluripotent Stem Cells. *J Cell Physiol*. 2015 Nov;230(11):2640-6.
11. Zhang Y, Chen ML, Zhou Y, Yi L, Gao YX, Ran L, Chen SH, Zhang T, Zhou X, Zou D, Wu B, Wu Y, Chang H, Zhu JD, Zhang QY, Mi MT. Resveratrol improves hepatic steatosis by inducing autophagy through the cAMP signaling pathway. *Mol Nutr Food Res*. 2015 Aug;59(8):1443-57.
12. Chen B, Lin T, Yang X, Li Y, Xie D, Cui H. Intermittent parathyroid

- hormone (1-34) application regulates cAMP-response element binding protein activity to promote the proliferation and osteogenic differentiation of bone mesenchymal stromal cells, via the cAMP/PKA signaling pathway. *Exp Ther Med*. 2016 Jun;11(6):2399-2406.
13. Jia B, Gao Y, Li M, Shi J, Peng Y, Du X, Klocker H, Sampson N, Shen Y, Liu M, Zhang J. GPR30 Promotes Prostate Stromal Cell Activation via Suppression of ER α Expression and Its Downstream Signaling Pathway. *Endocrinology*. 2016 Aug;157(8):3023-35.
 14. Shen C, Li X, Rasooly A, Guo L, Zhang K, Yang M. A single electrochemical biosensor for detecting the activity and inhibition of both protein kinase and alkaline phosphatase based on phosphate ions induced deposition of redox precipitates. *Biosens Bio electron*. 2016 Nov 15;85:220-5.
 15. Chen T, Xu DZ, Li Q, Mou P, Zeng Z, Brass LF, Zhu L. The regulation of Sema4D exodomain shedding by protein kinase A in platelets. *Platelets*. 2016 Nov;27(7):673-679.
 16. Yu ZL, Li DQ, Huang XY, Xing X, Yu RQ, Li Z, Li ZB. Lysophosphatidic acid upregulates connective tissue growth factor expression in osteoblasts through the GPCR/PKC and PKA pathways. *Int J Mol Med*. 2016 Feb;37(2):468-74.
 17. Deng X, Deng L, Wang P, Cheng C, Xu K. Post translational modification of CREB 1 decreases collagen I expression by inhibiting the TGF β 1 signaling pathway in rat hepatic stellate cells. *Mol Med Rep*. 2016 Dec;14(6):5751-5759.
 18. Shen XF, Teng Y, Sha KH, Wang XY, Yang XL, Guo XJ, Ren LB, Wang XY, Li J, Huang N. Dietary flavonoid luteolin attenuates uropathogenic Escherichia. Coli invasion of the urinary bladder. *Biofactors*. 2016 Nov 12;42(6):674-685.
 19. Xu Y, Zhao Y, Xu Y, Guan Y, Zhang X, Chen Y, Wu Q, Zhu G, Chen Y, Sun F, Wang J, Yu Y. Blocking inhibition to YAP by ActinomycinD enhances anti-tumor efficacy of Corosolic acid in treating liver cancer. *Cell Signal*. 2017 Jan;29:209-217.

Version 2016.12.09